Studies of the fast ion confinement in the Large Helical Device by using neutron measurement and integrated codes

journal or publication title	Journal of Plasma Physics
volume	86
number	3
page range	815860306
year	2020-06-17
URL	http://hdl.handle.net/10655/00012763

doi: 10.1017/S0022377820000525

Studies of the Fast Ion Confinement in LHD by Using Neutron Measurement and Integrated Codes

H. Nuga^{1†}, R. Seki^{1,2}, K. Ogawa^{1,2}, S. Kamio¹, Y. Fujiwara¹, M. Osakabe^{1,2}, M. Isobe^{1,2}, T. Nishitani¹, M. Yokoyama^{1,2}, and LHD Experiment Group¹

¹National Institute for Fusion Sciences, National Institutes of Natural Sciences, Toki, Japan

²The Graduate University for Advanced Studies, SOKENDAI, Toki, Japan

(Received xx; revised xx; accepted xx)

The neutral beam (NB) fast ion confinement in the Large Helical Device (LHD) is studied for several full field ($B_t \sim 2.75$ T) magnetic configurations by the combination between the neutron measurement and simulations. To investigate the NB fast ion confinement, we have performed a series of short-pulse NB injection experiments. The experiment results are analyzed by integrated code TASK3D-a. From this investigation, the effective particle diffusion coefficient of the tangential and perpendicular NBs are approximately $D^{\text{eff}} \sim 0.1\text{m}^2/\text{s}$ and $D^{\text{eff}} \sim 1\text{m}^2/\text{s}$ in the standard configuration. It is clarified that the NB fast ion confinement improves when the plasmas are shifted inward. Moreover, it is also found that the simulation, which considers the deuteron dilution effect due to the presence of impurity ions, can describe the neutron emission rate consistent to the measurement.

1. Introduction

Generally, the temperature of the fusion plasmas is sustained by the external heating. The heatings are achieved by the kinetic energy transfer from fast particles to the bulk plasmas through the Coulomb collision. Additionally, in the future fusion power plant, the fusion plasmas are mainly sustained by the fusion-born α particles, which have 3.5 MeV kinetic energy. Therefore, fast particles should be confined until their velocities are decelerated to the plasma thermal velocity. For this reason, the fast particle confinement is one of the most important issues (Heidbrink & Sadler 1994; Gorelenkov *et al.* 2014) for magnetically confined fusion devices.

There are several diagnostics systems to investigate the fast particle physics (Gorelenkov *et al.* 2014), such as neutral particle analyzer (NPA) (Medley & Roquemore 1998; Osakabe *et al.* 2006; Kamio *et al.* 2019), fast ion D_α spectroscopy (FIDA) (Luo *et al.* 2004; Heidbrink *et al.* 2007), fast ion loss detector (FILD) (Darrow *et al.* 1995; Ogawa *et al.* 2010), and neutron spectroscopy (Adams *et al.* 1993; Ishikawa *et al.* 2007). Because the deuterium plasma experiment campaign began in the Large Helical Device (LHD) (Takeiri 2018a,b; Osakabe *et al.* 2018, 2017) from March 2017, the investigation of fast ion behavior using neutron diagnostics (Isobe *et al.* 2018; Ogawa *et al.* 2019) and FIDA (Fujiwara *et al.* 2019) have been enabled in LHD.

In LHD deuterium plasmas, ~ 2.45 MeV fast neutrons are yielded owing to the deuterium-deuterium (DD) fusion reaction. Because of the high injection energy neutral

† Email address for correspondence: nuga.hideo@nifs.ac.jp

beams up to $E_b \sim 180$ keV, most of the neutrons are generated by the fusion reaction between fast deuteron and thermal deuteron instead of the reaction between thermal ions (Osakabe *et al.* 2017). This is the so-called “beam-thermal” fusion reaction. In such cases, we can study the fast ion global confinement from the neutron emission rate S_n . This is because the DD fusion reactivity increases monotonically with their relative kinetic energy up to $E \sim 2$ MeV. If the plasma temperature and density are constant in time, the decrement of the neutron emission rate comes from the decrement of the fusion cross-section due to the fast ion deceleration and the decrement of the fast ion density. Because the fast ion deceleration can be estimated by simulations, we can estimate the fast ion confinement from the neutron measurement without simulating the fast ion transport, which is usually too complex to estimate accurately.

We have studied the effective confinement time of NB fast ions in LHD using neutron measurement in this paper. There are several previous works to investigate fast ion confinement through the neutron diagnostics. Earliest works in tokamaks (Strachan *et al.* 1981; Heidbrink *et al.* 1988, 1991) showed that the classical collisional slowing down model can explain the neutron emission within good agreement. The following investigation in JT-60U tokamak (Tobita *et al.* 1994) reported the deviation from the classical collisional slowing down model due to the toroidal field ripple for perpendicular NB. Anomalous fast ion losses, which can not be explained by the toroidal field ripple, were observed on TFTR (Ruskov *et al.* 1999). Moreover, fast ion confinement was investigated by using neutron measurement also in various types of magnetically confinement device such as the reversed field pinch devices (Fiksel *et al.* 2005; Anderson *et al.* 2014), the spherical tokamak (Heidbrink *et al.* 2003), the tokamak (Zhang *et al.* 2012), and the stellarator and heliotron (Isobe *et al.* 2001; Nuga *et al.* 2019). In our previous work (Nuga *et al.* 2019), we have reported the NB fast ion confinement time by using neutron measurement for single magnetic configuration in LHD. In this paper, we investigate the dependence of the fast ion confinement on the magnetic configuration and discuss the contribution of the charge exchange loss to the fast ion confinement especially in low electron density plasmas.

The rest of this paper consists of the following. The simple instruction of the experimental scenario and experiment devices are provided in section 2. Simulation tools and models used in this paper are introduced in section 3. The results of the analyses to estimate the fast ion confinement time are shown in section 4 and section 5. The dilution effect of the deuteron density due to the presence of impurity ions is discussed in section 6. Section 7 concludes this paper.

2. Experimental setup

2.1. Experimental scenario

A series of experiments which use short pulse NB injection have been performed in LHD to investigate the fast ion confinement time. The decay time of S_n , yielded by DD fusion reaction has been observed after short pulse NBs have been turned off. In the previous work (Nuga *et al.* 2019), we performed the short pulse NB experiment in single magnetic configuration ($R_{ax} = 3.6$ m, $B_t = 2.75$ T), where R_{ax} and B_t indicate the position of the magnetic axis and the strength of the magnetic field on the magnetic axis. The direction of the magnetic field is counter clockwise in the top view. In this paper, we have performed the short pulse NB experiment in five magnetic configurations, (R_{ax}, B_t) = (3.53m, 2.80T), (3.55m, 2.79T), (3.60m, 2.75T), (3.75m, 2.64T), and (3.90m,

FIGURE 1. Typical waveform of this series of experiments (SN148442) is displayed. the NB port through power in each beam, the plasma temperature and density on the position, where the normalized minor radius $\rho = 0.5$, and the neutron emission rate are shown.

2.54T), respectively. Typical waveform of this series of experiments is shown in figure 1. NB pulse width is ~ 40 ms in each beam.

In this series of experiments, the fast ion confinement time, τ_c , is assumed to be:

$$\tau_c^{-1} = \tau_n^{-1} - (\tau_n^{\text{cl}})^{-1}, \quad (2.1)$$

where τ_n is the neutron decay time after NB has been turned off and τ_n^{cl} is the e -folding time of the neutron decay rate due to the classical fast ion deceleration. This is because the reduction of the neutron emission rate is caused by the decrease of the fusion cross-section due to the deceleration and the decrease of the fast ion density due to the fast ion loss. Because the contribution of the fast ion loss is ignored at the estimation of τ_n^{cl} , τ_c can be estimated from τ_n and τ_n^{cl} . In this paper, τ_n can be obtained from the experimental results and τ_n^{cl} can be obtained from the simulation results instead of the analytical solution given by the reference (Strachan *et al.* 1981). The method of the simulation will be introduced in section 3.

2.2. Experiment Apparatus

In this series of experiments, two external heating systems, electron cyclotron heating (ECH) and NBI, are used. Plasmas are sustained by 3.5 MW ECH.

The top view of the NBI system in LHD (Takeiri *et al.* 2010) is shown in figure 2. Typical injection energy and power in each beam is also displayed. In LHD, there are three negative ion-based tangential NBs and two positive ion-based perpendicular NBs. The direction of NB#1 and NB#3 are co-direction to the magnetic field and the direction of NB#2 is counter-direction to the magnetic field. It is noted that the injection energy of NB#4 is set to be untypically low energy, $E_b \sim 35$ keV, to investigate the dependence of the perpendicular NBs on the injection energy.

Experiments have been performed with five magnetic configurations as noted in sec. 2.1. The averaged plasma minor radius, which depends on the position of the magnetic axis, is approximately $0.58 - 0.63$ m.

LHD is equipped with several diagnostic systems (Kawahata *et al.* 2010). In this paper, we have mainly used electron temperature and density profiles, the ion ratio among proton, deuteron, and helium ion, and the time evolution of the neutron emission rate. The Thomson scattering diagnostics is used for the measurement of the electron density and temperature (Narihara *et al.* 2001; Yamada *et al.* 2010). A fast-response, wide

FIGURE 2. Top view of NBI system in LHD. There are three tangential beams, NB#1, NB#2, and NB#3, and there are two perpendicular beams, NB#4 and NB#5. Because the direction of the magnetic field line is counter-clockwise, NB#1 and NB#3 is the co-direction to the magnetic field line.

FIGURE 3. Flow of the simulations to obtain the time evolution of the neutron emission rate.

dynamics range Neutron Flux Monitor (NFM) (Isobe *et al.* 2018) is equipped in LHD to measure the neutron emission rate.

3. Simulation tools

Estimation of τ_n^{cl} is required to evaluate τ_c as shown in eq. 2.1. Although there is the analytical expression of τ_n^{cl} (Strachan *et al.* 1981), we use the following simulations to include the plasma profiles, plasma volume, and NB birth profile for the estimation of τ_n^{cl} .

In this paper, the neutron emission simulations have been performed by the combination among VMEC (Hirshman & Whitson 1983), FIT3D (HFREYA and

MCNBI)(Murakami *et al.* 1995; Sato *et al.* 2008; Vincenzi *et al.* 2016; Seki *et al.* 2019), and CONV_FIT3D(Nuga *et al.* 2019), which are included in the integrated transport analysis suite TASK3D-a(Yokoyama *et al.* 2017). The flow of the simulations are displayed in figure 3. Three-dimensional magnetic equilibrium is calculated by VMEC code. According to the obtained magnetic configuration, FIT3D calculates NB fast ion birth profile. At this step, NB fast ion orbits are followed for a few tens of micro seconds by using Monte Carlo method to consider the NB fast ion prompt loss effect. In this process, the effect of the birth profile broadening owing to the deviation of the NB fast ion drift orbit from the magnetic surface is also considered. Moreover, for positive ion-based NBs, the full, half, and one third energy fractions of injected neutral particles are assumed to be 0.78, 0.16, and 0.06, respectively. Although these values are the typical values of perpendicular NBI systems in LHD, it is not reliable for NB#4. This is because NB#4 has untypically low injection energy in the following analyses. CONV_FIT3D solves the NB fast ion slowing down equation to obtain the energy distribution of NB fast ion. Here, the fast ion transport across the magnetic surface during slowing down process is neglected. In LHD, the beam-thermal fusion reaction is dominant instead of the reaction between thermal ions due to the insufficient plasma temperature. Additionally, in this paper, the fusion reaction between NB fast ions is also negligible. Therefore, it is enough to consider the fusion reaction between NB fast ion and thermal ion in this paper. Because it can be considered that the velocity distribution of thermal ion is isotropic, the velocity direction of NB fast ion is not necessary for the estimation of the fusion reaction rate. For this reason, CONV_FIT3D, which can obtain the energy distribution of NB fast ion, can estimate the neutron emission rate. The details of CONV_FIT3D are introduced in section 3.1.

3.1. CONV_FIT3D

CONV_FIT3D is a code solving the fast ion slowing down equation based on the classical Coulomb collision theory. In the previous work(Nuga *et al.* 2019), we had chosen the model which assumes NB fast ion velocity v_{EP} is sufficiently faster than thermal ion velocity v_{th} . However, in this paper, we have a perpendicular NB#4, which has an injection energy of $E_b = 35$ keV. NB fast ions, whose kinetic energy is 35 keV, can not satisfy the assumption $v_{EP} \gg v_{th}$ because typical temperature of this series of experiments is a few keV. Therefore, we have refined the slowing down equation based on the eq. (30c) in reference (Karney 1986). This model assumes the background Maxwellian. The refined equations are expressed as below:

$$\frac{dE}{dt} = -\frac{2}{\tau_{se}} Z_{EP}^2 E \times \left[1 + \frac{3\pi^{1/2} m_{EP}^{3/2} T_e^{3/2}}{4m_e^{1/2}} \left(\sum_i \frac{n_i Z_i^2}{n_e m_i} g(u_i) \right) \frac{1}{E^{3/2}} \right] \quad (3.1)$$

$$g(u_i) = \text{erf}(u_i) - u_i \text{erf}'(u_i) \quad (3.2)$$

$$\tau_{se} = \frac{3(2\pi)^{3/2} \epsilon_0^2 m_{EP} T_e^{3/2}}{n_e e^4 m_e^{1/2} \ln \Lambda} \quad (3.3)$$

$$\text{erf}(u) = \frac{2}{\pi} \int_0^u \exp(-x^2) dx \quad (3.4)$$

$$\text{erf}'(u) = \frac{2}{\pi} \exp(-u^2), \quad (3.5)$$

FIGURE 4. Example of the neutron emission simulation assuming $n_D = n_e$ and $\tau_c = \infty$ is displayed.

where τ_{se} is the Spitzer slowing down time on electron, $u_i = v/\sqrt{2}v_{th,i}$ and $v_{th} = \sqrt{T_i/m_i}$. In the following calculation, we assumed that the ion temperature T_i is equal to the electron temperature T_e . In actuality, however, it is expected that T_e is greater than T_i because the plasmas are sustained by ECH.

As noted above, in LHD, the beam-thermal fusion reaction is dominant. In this case, the fusion reaction rate can be expressed as:

$$\mathcal{R}_{bt} = n_D \int \langle \sigma v \rangle_{bt}(T_i, E_{EP}) f_{EP}(\mathbf{v}_{EP}) d\mathbf{v}_{EP}, \quad (3.6)$$

where n_D is the deuteron density in the bulk plasma. The beam-thermal reactivity $\langle \sigma v \rangle_{bt}$, which is derived by Mikkelsen(Mikkelsen 1989), is implemented in CONV_FIT3D. The fitting coefficients used in the expression of $\langle \sigma v \rangle_{bt}$ are chosen from reference(Bosch & Hale 1993).

4. Analysis results for tangential beams

In figure 4, the example of the simulated neutron emission rate is shown with the measured neutron emission rate. The solid curve is the measured neutron emission rate and the dashed curve is the simulated neutron emission rate. The simulation overestimates the neutron emission rate due to the assumptions that $n_D = n_e$ and $\tau_c = \infty$. From measured and simulated neutron emission rates, we can estimate τ_n and τ_n^{cl} by using the least mean square fitting. The fitting functions are $f(t) = S_n^{p,exp} \exp(-t/\tau_n)$ for measured decay rate and $f(t) = S_n^{p,sim} \exp(-t/\tau_n^{cl})$ for simulated decay rate, where $S_n^{p,exp}$ and $S_n^{p,sim}$ are the peak values of the measured and simulated neutron emission rate. The durations which is used for this fitting are defined as the times when the neutron emission rates decay to $S_n^{p,exp}/e$ and $S_n^{p,sim}/e$, respectively.

Similar experiments have been performed with the same magnetic configuration ($R_{ax} = 3.53$ m, $B_t = 2.8045$ T) and the different plasma parameters to survey the dependence of τ_n on τ_n^{cl} . It is noted that the control parameter is the electron density n_e in these experiments. Additionally, because the plasma heating power is constant in each shot, the electron temperature T_e also changes with the electron density. Figure 5 is obtained from experiment and simulation results for three tangential NBs. Their injection energies are also displayed in fig. 5. In this figure, x axis denotes the simulated neutron decay time τ_n^{cl} and y axis denotes the experimental neutron decay time τ_n . The y error-bars shown in fig. 5 denote the standard deviation of τ_n . The solid guideline indicates $y = x$. If the NB fast

FIGURE 5. The experimentally measured neutron decay times are plotted against the simulated neutron decay time for tangential NBs in the configuration of $R_{\text{ax}} = 3.53$ m. Dashed guide curve is $\tau_n = \tau_c \tau_n^{\text{cl}} / (\tau_n^{\text{cl}} + \tau_c)$, $\tau_c = 0.53$ s.

ions are confined perfectly, plots obey this guideline. Actually, because the NB fast ion loss is not ignorable, plots separate from the guide line as shown in fig. 5. Additionally, in fig. 5, there is the dashed guide curve which indicates $\tau_n = 0.53 \tau_n^{\text{cl}} / (\tau_n^{\text{cl}} + 0.53)$. The reason for the value $\tau_c = 0.53$ is explained later. From fig. 5, it is found that the different trend, which does not obey the dashed guide curve, appears in the $\tau_n^{\text{cl}} > 0.4$ s region. This means that the dominant transport mechanism is different between the $\tau_n^{\text{cl}} < 0.4$ s region and the $\tau_n^{\text{cl}} > 0.4$ s region.

It is noted that high frequency (> 10 kHz) MHD instabilities triggered by NBI hardly appeared in short τ_n^{cl} region. On the contrary, in long τ_n^{cl} region, NB fast ion driven instabilities were observed in several discharges. Therefore, we can ignore the anomalous transport due to fast ion driven instabilities in short τ_n^{cl} region. The following analyses in long τ_n^{cl} region may include the anomalous transport. The low frequency (< 10 kHz) MHD instabilities excited by bulk plasma pressure were observed through this series of experiments. The typical amplitude of the low frequency magnetic fluctuation, which is detected by magnetic probe set on the vacuum vessel, were $\delta B/B_t < 2 \times 10^{-5}$. This value is sufficiently low as compared to the case that the low frequency magnetic fluctuation plays important role to the NB fast ion loss (Ogawa *et al.* 2010). Therefore, it can be considered that NB fast ion loss due to the low frequency fluctuation can be negligible.

4.1. Long τ_n^{cl} region ($\tau_n^{\text{cl}} > 0.4$ s)

In LHD plasmas, it is known that the neutral atom density in plasma core region has negative n_e dependence (Fujii *et al.* 2015; Goto *et al.* 2011, 2016) because the penetration of neutral atoms into the plasma core increases with the decrease of the plasma density. Therefore, it can be considered that the neutral atom density in plasma core region in $\tau_n^{\text{cl}} > 0.4$ sec region is higher than that in $\tau_n^{\text{cl}} < 0.4$ sec region. Here we focus on the $\tau_n^{\text{cl}} > 0.4$ s region to investigate the contribution of the charge exchange loss to the fast ion loss.

From fig. 5, it is found that the values of τ_n for NB#1, NB#2, and NB#3 are approximately $\tau_n \sim 0.25$ s, 0.2 s, and 0.2 s at $\tau_n^{\text{cl}} = 1$ s. Here we assume that the separation from the dashed guide curve in long τ_n^{cl} region can be described only by the charge exchange loss. In this assumption, the neutron decay time τ_n can be expressed as:

$$\tau_n^{-1} = (\tau_n^{\text{cl}})^{-1} + (\tau_c^{\text{eff}})^{-1} + \tau_{\text{cx}}^{-1}, \quad (4.1)$$

where τ_c^{eff} is the effective confinement time in short τ_n^{cl} region and τ_{cx} is the charge exchange time. In this configuration, the value of τ_c^{eff} , which will be obtained in section

FIGURE 6. The dependence of the inverse of the charge exchange reactivity $1/\langle\sigma v\rangle_{cx}$ on beam energy is shown. Here, it is assumed that the velocity of neutral atom is negligible.

FIGURE 7. The short τ_n^{cl} region ($\tau_n^{cl} < 0.4$ s) of figure 5 is focused.

4.2, is $\tau_c^{eff} = 0.53$ s. For these assumptions and estimations, the charge exchange times for NB#1, NB#2, and NB#3 are approximately $\tau_{cx} = 0.90$ s, 0.47 s, and 0.47 s, respectively.

The dependence of the charge exchange reactivity on the beam energy is shown in fig. 6. Although fast ions in plasmas have a velocity distribution, we estimate the charge exchange reactivity $\langle\sigma v\rangle_{cx}$ in each beam from the beam injection energy. This is because that we have discussed the fast ion confinement through the measurement of the neutron emission rate, which exponentially depends on the beam energy within the range of NB injection energy. Accordingly, the charge exchange time can be expressed as $\tau_{cx}^{-1} = n_0 \langle\sigma v\rangle_{cx}(E_b)$, where n_0 is the neutral atom density. From this expression, we can estimate the values of n_0 in each beam. The neutral atom densities, which are felt by fast ions from NB#1, NB#2, and NB#3, are $n_0 = 1.7 \times 10^{14} \text{m}^{-3}$, $1.9 \times 10^{14} \text{m}^{-3}$, and $2.1 \times 10^{14} \text{m}^{-3}$, respectively. In the reference (Goto *et al.* 2016), the neutral atom density in the LHD plasma, whose electron density is $\sim 2 \times 10^{19} \text{m}^{-3}$, is measured as approximately 10^{14}m^{-3} . Because the plasmas in long τ_n^{cl} region have the electron density of approximately $n_e \lesssim 0.5 \times 10^{19} \text{m}^{-3}$, it can be expected that the neutral density is larger than 10^{14}m^{-3} in these plasmas. Therefore, it can be considered that the estimation of n_0 is within the realistic value and the charge exchange loss is not negligible in long τ_n^{cl} region.

4.2. Short τ_n^{cl} region ($\tau_n^{cl} < 0.4$ s)

Here we focus on the region $\tau_n^{cl} < 0.4$ s of fig. 5 to exclude the cases that the charge exchange loss is significant. The dashed guide curve in figure 7 is obtained by the weighted least mean square fitting in $\tau_n = \tau_n^{cl} \tau_c^{eff} / (\tau_n^{cl} + \tau_c^{eff})$ function in $\tau_n^{cl} < 0.4$ s region. It is noted that because three tangential NBs have a similar trend, they are fitted in a single

FIGURE 8. The effective particle diffusion coefficients are plotted against the position of the magnetic axis.

FIGURE 9. The experimentally measured neutron decay times are plotted against the simulated neutron decay time for perpendicular NBs in the configuration of $R_{ax} = 3.53$ m. Dashed guide curve is $\tau_n = \tau_c \tau_n^{cl} / (\tau_n^{cl} + \tau_c)$, $\tau_c = 0.057$ s.

curve. For this fitting, the effective fast confinement time in this region is obtained as $\tau_c^{eff} = 0.53$ s.

Similar analyses are performed for different configurations shown in section 2.1. To compare the confinement among different magnetic configurations, the effective fast ion confinement time τ_c^{eff} is converted into the effective particle diffusion coefficient D^{eff} . The relation between τ_c^{eff} and D^{eff} is expressed as

$$D^{eff} = \frac{a^2}{5.8\tau_c^{eff}}, \quad (4.2)$$

where a is the plasma minor radius. Here it is assumed that D^{eff} is uniform in the plasma and the plasma poloidal cross-section is not elongated. The R_{ax} dependence of D^{eff} is shown in figure 8. It is found that the fast ion confinement becomes poor when the plasma is shifted outward. Additionally, the absolute value of D^{eff} obtained in this paper is larger than that obtained in tokamaks (Tobita *et al.* 1994; Heidbrink *et al.* 1988).

5. Analysis Results for perpendicular beams

Perpendicular beams are analyzed by a method similar to the tangential beams. Figure 9 shows the relation between τ_n and τ_n^{cl} for perpendicular NBs. The dashed guide curve indicates $\tau_n = 0.057\tau_n^{cl} / (\tau_n^{cl} + 0.057)$. In figure 9, the different trends appear at approximately $\tau_n^{cl} = 0.1$ sec.

FIGURE 10. The short τ_n^{cl} region ($\tau_n^{\text{cl}} < 0.1$ s) of figure 9 is focused on.

FIGURE 11. The effective particle diffusion coefficients for perpendicular beams are plotted against the position of the magnetic axis.

5.1. Long τ_n^{cl} region ($\tau_n^{\text{cl}} > 0.1$ s)

Similar to section 4.1, we investigate the contribution of the charge exchange loss to the fast ion confinement. From fig. 9, the neutron decay time of NB#4 and NB#5 at $\tau_n^{\text{cl}} = 0.2$ s are approximately $\tau_n = 0.025$ s and 0.035 s, respectively. From these values and eq. (4.1), the charge exchange times of NB#4 and NB#5 can be estimated as $\tau_{\text{cx}} = 0.057$ s and 0.17 s, respectively. The neutral atom densities evaluated from the τ_{cx} are $n_0 = 2.1 \times 10^{14} \text{m}^{-3}$, and $1.3 \times 10^{14} \text{m}^{-3}$, respectively. These values of density are within the range of realistic value as noted in sec. 4.2. Therefore, it can be also considered that the charge exchange loss are not negligible for perpendicular beams in long τ_n^{cl} region.

5.2. Short τ_n^{cl} region ($\tau_n^{\text{cl}} < 0.1$ s)

Owing to the above discussion, we focus on the $\tau_n^{\text{cl}} < 0.1$ sec region to exclude the cases that the charge exchange loss is significant. In figure 10, the short τ_n^{cl} region $\tau_n^{\text{cl}} < 0.1$ sec of figure 9 is focused on. The guide curve indicates $\tau_c^{\text{eff}} \tau_n^{\text{cl}} / (\tau_n^{\text{cl}} + \tau_c^{\text{eff}})$, $\tau_c^{\text{eff}} = 0.057$ sec. The value of τ_c^{eff} is obtained by the weighted least mean square fitting in $\tau_n = \tau_c^{\text{eff}} \tau_n^{\text{cl}} / (\tau_n^{\text{cl}} + \tau_c^{\text{eff}})$.

Similar to the discussion in section 4.2, the effective particle diffusion coefficient D^{eff} for perpendicular beam is estimated for the five magnetic configurations. In figure 11, the dependence of D^{eff} on R_{ax} for perpendicular beams are shown. It is found that the perpendicular beams have a poorer confinement than the tangential beams. The same as tangential beams, The confinement of the perpendicular beams becomes poor when the plasmas are shifted outward.

It is noted that the perpendicular beam injection energy is not uniform in each configuration due to experimental constraints. In the cases of $R_{\text{ax}} = 3.75$ m and 3.9 m, the perpendicular beam injection energy is $E_b = 60$ keV. The data of $R_{\text{ax}} = 3.6$ m configuration consist of the data with $E_b = 35$ keV, 60 keV, and 80 keV. In the remaining

FIGURE 12. The relation between $S_n^{p,exp}$ and $\mathcal{F}S_n^{p,sim}$ is shown for the $R_{ax} = 3.53$ m configuration. The solid line indicates $y = x$ and the dashed line indicates $y = 0.42x$.

configurations ($R_{ax} = 3.53$ m and 3.55 m), the data consist of those with $E_b = 35$ keV and 80 keV. These data are fitted in a single curve as shown in fig. 10.

6. Discussion regarding the deuteron dilution

In this paper, in order to estimate the fast ion confinement time, we have focused on the neutron decay time instead of the absolute value of the neutron emission rate itself. This is because that there are some ambiguities for estimating the absolute value of the neutron emission rate. The fusion reaction rate in beam-thermal reaction dominant plasmas is described as eq. 3.6. Our simulation over-estimates the neutron emission rate due to the assumptions of $n_D = n_e$ and $\tau_c = \infty$ as shown in fig. 4. In actual plasmas, however, deuterons are diluted due to the presence of impurity ions. Because the evaluation of the impurity density profile is difficult, the ambiguity can not be reduced to estimate the neutron emission rate. On the other hand, the estimation of the neutron decay time has fewer ambiguities. This is because the fast ion slowing down process mainly depends on the measurable values, which are electron density n_e and temperature T_e . Since we have estimated the fast ion confinement time τ_c until the previous section, the diluted deuteron density can be evaluated by the comparison between the peak values of measured and simulated neutron emission rate.

The relation between two neutron emission rate peak values, measured peak value $S_n^{p,exp}$ and simulated peak value $S_n^{p,sim}$, can be expressed as (Nuga *et al.* 2019):

$$\frac{S_n^{p,exp}}{S_n^{p,sim}} = \frac{n_D}{n_e} \mathcal{F},$$

$$\mathcal{F} \equiv \frac{\tau_c}{\tau_n^{cl} + \tau_c} \frac{1 - \exp(-t_0/[\tau_n^{cl}\tau_c/(\tau_n^{cl} + \tau_c)])}{1 - \exp(-t_0/\tau_n^{cl})}, \quad (6.1)$$

where t_0 is the beam injection duration $t_0 \sim 40$ ms. Here we assume that the plasma parameters and the fast ion birth rate are constant during beam injection duration.

Figure 12 shows the relation between $S_n^{p,exp}$ and $\mathcal{F}S_n^{p,sim}$ for the $R_{ax} = 3.53$ m configuration. It is noted that, in this figure, data in long τ_n^{cl} region ($\tau_n^{cl} > 0.4$ s for tangential beams and $\tau_n^{cl} > 0.1$ s for perpendicular beams) are omitted. From eq. (6.1), the slope of dashed line in fig. 12 indicates the ratio of the deuteron density to the electron density n_D/n_e . The value is obtained by the weighted least mean square fitting in $y = \alpha x$, where α is the fitting parameter. For this reason, here we assume the deuteron density ratio to the electron density is approximately $n_D/n_e = 0.42$.

It is noted that this fitting is performed by using only tangential NBs. This is because the full energy fraction of perpendicular NBs is unclear. From fig. 12, plots of NB#5

FIGURE 13. The carbon density ratio to the electron density n_C/n_e is shown for three discharges. The horizontal axis indicates the normalized minor radius ρ . The minus value of ρ means the position inside the magnetic axis.

has a good agreement to the fitting line. This indicate that the assumption of the full energy fraction, 0.78, used in the simulation is valid. On the contrary, results of NB#4 are separated from the fitting line. One of the reasons of this deviation is that the assumption of the full energy fraction is invalid. If the deviation can be explained only by the full energy fraction, the fraction becomes 1.2. Therefore, it seems that there is the other factor separating the results from the fitting. In this paper, NB#4 has a low injection energy, whose value is out of typical range of injection energy. Because of this unusual operation of NB#4, the S_n peak value estimation of NB#4 is not reliable.

The deuterium ion ratio and the helium ion ratio in this series of experiment are approximately $n_D/(n_H + n_D) \sim 1$ and $(n_D + n_H)/(n_D + n_H + n_{He}) \sim 0.9$. Here, we assume that there are only three ion species, deuterium, helium ion, and carbon ion in plasmas. This assumption indicates $n_e = n_D + 2n_{He} + 6n_C$. We can estimate the typical ion ratios from the deuterium ratio estimation, two measured ratios, and this assumption. These ratios are $n_D/n_e = 0.42$, $n_{He}/n_e = 0.047$, and $n_C/n_e = 0.081$. The effective charge of the plasma, which consist of these ion ratios, is approximately $Z_{eff} = 3.5$. From references (Zhou *et al.* 2010; Huang *et al.* 2015), the typical value of the effective charge in LHD plasmas measured by EUV spectrometer is approximately $Z_{eff} = 2 - 4$. Therefore, this estimation is within the range of the realistic value.

Additionally, the carbon density profile, n_C , is measured by the charge exchange spectroscopy of CVI using neutral beam (Chen *et al.* 2019; IDA *et al.* 2019) in LHD. Unlike $n_D/(n_H + n_D)$ and $(n_D + n_H)/(n_D + n_H + n_{He})$, the carbon density ratio to the electron density changes widely shot-by-shot as shown in figure 13. In most of the discharges, the value of n_C/n_e has changed around $\lesssim 0.1$ and it does not exceed 0.2. Therefore, it can be considered that the value $n_D/n_e = 0.42$ is reasonable.

Here we can recalculate the neutron emission rate by using estimated τ_c and ion ratios. Figure 14 shows the time evolutions of the measured and simulated neutron emission rate. In this simulation, the dilution effect of deuterium density and the fast ion loss effect are included. As compared to fig. 4, it is found that the consistent result for the measured neutron emission rate can be obtained by this simulation.

7. Conclusion

We have investigated the NB fast ion confinement in LHD deuterium plasmas in five magnetic configurations. The effective confinement time of NB fast ion has been estimated from the decay time of the neutron emission rate after short pulse NBs are turned off. It can be considered that the decay of the neutron emission rate can be expressed by the combination between the reduction of the fusion cross-section due to the fast ion slowing

FIGURE 14. Time evolutions of the measured and simulated neutron emission rate are shown. The bulk deuteron density is assumed to be $n_D/n_e = 0.42$ and the fast ion confinement times are assumed to be $\tau_c = 0.6$ s for tangential beams and $\tau_c = 0.06$ s for perpendicular beams.

down and the reduction of the fast ion density due to the fast ion loss. Therefore, the comparison between the simulated neutron decay time τ_n^{cl} , which ignores the fast ion loss, and the measured neutron decay time τ_n clarifies the fast ion confinement time.

As shown in figure 5 and figure 9, the relation between τ_n and τ_n^{cl} has two trends. In short τ_n^{cl} region, τ_n obeys the curve, $f(\tau_n^{\text{cl}}) = \tau_c \tau_n^{\text{cl}} / (\tau_n^{\text{cl}} + \tau_c)$, where τ_c is constant. In other words, this region is the high electron density region. On the other hand, τ_n separates from the guide curve as τ_n^{cl} increases. This indicates that the dominant fast ion loss mechanism is different in each region. As shown in section 4 and section 5, in long τ_n^{cl} region, the charge exchange loss can not be negligible because the low electron density plasmas include many neutral particles. To reduce the charge exchange loss in the estimation of τ_c , we have focused on the short τ_n^{cl} region. Here the short τ_n^{cl} region indicate $\tau_n^{\text{cl}} < 0.4$ sec for tangential beams and $\tau_n^{\text{cl}} < 0.1$ sec for perpendicular beams, respectively. In this region, we can estimate the effective confinement time τ_c^{eff} by the weighted least mean square fitting in $f(\tau_n^{\text{cl}})$. In addition, similar analyses have been performed for the other magnetic configurations. In figures 8 and 11, the effective NB fast ion diffusion coefficient D^{eff} is plotted against the position of the magnetic axis. It is clear that the tangential NBs have good confinement rather than the perpendicular NBs. Additionally, inward shifted plasmas have a good confinement of NB fast ions in both injection directions.

The dependence of the neo-classical diffusion coefficient on the position of magnetic axis in LHD have been discussed numerically in the reference (Murakami *et al.* 2002). This reference showed that the neo-classical diffusion coefficient in $1/\nu$ regime decreases as the plasma shifting inward. Because the neo-classical diffusion is dominated by trapped particles in usual, it can be considered that the dependence of D^{eff} on R_{ax} for perpendicular beam ions has same trend to the reference result. For tangential beam ions, it can be roughly explained that the orbit deviation from the magnetic surface becomes larger as the plasma shifting outward because the toroidal curvature component of the magnetic field, $B_{1,0}$ (poloidal mode number = 1, toroidal mode number = 0), increases as the plasma shifting outward. Obviously, however, the sensitivity of D^{eff} to R_{ax} is different from the reference results. Since the detailed discussion of this issue requires orbit simulations, this is the future work.

We have focused on the decay time of the neutron emission rate to estimate τ_c^{eff} instead of the absolute value of the neutron emission rate S_n . The absolute value of S_n linearly depends on the bulk deuteron density, which is difficult to measure in the plasma core region. In actual plasmas, the deuteron density is diluted by the presence of the impurity ions. This dilution effect can be estimated by the obtained τ_c^{eff} and by the difference of

the peak value of the measured and simulated neutron emission rate. From figure 12, the ratio of the deuteron density to the electron density is roughly estimated as $n_D/n_e = 0.42$. This estimation has no significant contradictions with impurity measurements as shown in section 6. According to the obtained τ_c and n_D/n_e , we can simulate the consistent neutron emission rate to the measured neutron emission rate.

Although we have estimated quantitatively the fast ion confinement time τ_c in LHD plasmas, the transport mechanism itself is still unclear. The clarification of the transport mechanism dominant in the short τ_n^{cl} region is the future work.

Acknowledgments

This work is supported partly by LHD project budgets (NIFS07KLPH004 and NIFS14KNTT025). It is performed on “Plasma Simulator” (FUJITSU FX100) of NIFS with the support and under the auspices of the NIFS Collaboration Research program (NIFS18KNSR007).

REFERENCES

- ADAMS, J. M., JARVIS, O. N., SADLER, G. J., SYME, D. B. & WATKINS, N. 1993 The jet neutron emission profile monitor. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **329** (1-2), 277–290.
- ANDERSON, J. K., CAPECCHI, W., EILERMAN, S., KOLINER, J. J., NORNBERG, M. D., REUSCH, J. A., SARFF, J. S. & LIN, L. 2014 Fast ion confinement in the three-dimensional helical reversed-field pinch. *Plasma Physics and Controlled Fusion* **56** (9), 094006.
- BOSCH, H.-S. & HALE, G. M. 1993 Improved formulas for fusion cross-sections and thermal reactivities. *Nuclear Fusion* **33** (12), 1919.
- CHEN, J., IDA, K., YOSHINUMA, M., MURAKAMI, I., KOBAYASHI, T., YE, M. Y. & LYU, B. 2019 Effect of energy dependent cross-section on flow velocity measurements with charge exchange spectroscopy in magnetized plasma. *Physics Letters A* **383** (12), 1293–1299.
- DARROW, D. S., HERRMANN, H. W., JOHNSON, D. W., MARSALA, R. J., PALLADINO, R. W., ZWEBEN, S. J. & TUSZEWSKI, M. 1995 Measurement of loss of dt fusion products using scintillator detectors in tfr. *Review of scientific instruments* **66** (1), 476–482.
- FIKSEL, G., HUDSON, B., DEN HARTOG, D. J., MAGEE, R. M., O’CONNELL, R., PRAGER, S. C., BEKLEMISHEV, A. D., DAVYDENKO, V. I., IVANOV, A. A. & TSIDULKO, YU. A. 2005 Observation of weak impact of a stochastic magnetic field on fast-ion confinement. *Phys. Rev. Lett.* **95**, 125001.
- FUJII, K., GOTO, M., MORITA, S. & LHD EXPERIMENT GROUP 2015 Study of neutral hydrogen transport in lhd core plasmas based on high dynamic-range balmer- α spectroscopy. *Nuclear Fusion* **55** (6), 063029.
- FUJIWARA, Y., KAMIO, S., YAMAGUCHI, H., GARCIA, A. V., STAGNER, L., NUGA, H., SEKI, R., OGAWA, K., ISOBE, M., YOKOYAMA, M., NISHITANI, T., HEIDBRINK, W. W., OSAKABE, M. & LHD EXPERIMENT GROUP 2019 Evaluation of an energetic particle profile using a tangential-fida diagnostic in the large helical device. *Plasma and Fusion Research* **14**, 3402129–3402129.
- GORELENKOV, N. N., PINCHES, S. D. & TOI, K. 2014 Energetic particle physics in fusion research in preparation for burning plasma experiments. *Nuclear Fusion* **54** (12), 125001.
- GOTO, M., SAWADA, K., FUJII, K., HASUO, M. & MORITA, S. 2011 Evaluation of hydrogen atom density in the plasma core region based on the balmer- α line profile. *Nuclear Fusion* **51** (2), 023005.
- GOTO, M., SAWADA, K., OISHI, T. & MORITA, S. 2016 Particle source and edge confinement

- study based on spectroscopic diagnosis in the lhd. *Plasma Physics and Controlled Fusion* **58** (8), 084001.
- HEIDBRINK, W. W., BARNES, C. W., HAMMETT, G. W., KUSAMA, Y., SCOTT, S. D., ZARNSTORFF, M. C., JOHNSON, L. C., MCCUNE, D., MEDLEY, S. S., PARK, H. K., ROQUEMORE, A. L., STRACHAN, J. D. & TAYLOR, G. 1991 The diffusion of fast ions in ohmic tfrt discharges. *Physics of Fluids B: Plasma Physics* **3** (11), 3167–3170.
- HEIDBRINK, W. W., KIM, J. & GROEBNER, R. J. 1988 Comparison of experimental and theoretical fast ion slowing-down times in diii-d. *Nuclear fusion* **28** (10), 1897.
- HEIDBRINK, W. W., LUO, Y., BURRELL, K. H., HARVEY, R. W., PINSKER, R. I. & RUSKOV, E. 2007 Measurements of fast-ion acceleration at cyclotron harmonics using balmer-alpha spectroscopy. *Plasma Physics and Controlled Fusion* **49** (9), 1457.
- HEIDBRINK, W. W., MIAH, M., DARROW, D., LEBLANC, B., MEDLEY, S. S., ROQUEMORE, A. L. & CECIL, F. E. 2003 The confinement of dilute populations of beam ions in the national spherical torus experiment. *Nuclear fusion* **43** (9), 883.
- HEIDBRINK, W. W. & SADLER, G. J. 1994 The behaviour of fast ions in tokamak experiments. *Nuclear Fusion* **34** (4), 535.
- HIRSHMAN, S. P. & WHITSON, J. C. 1983 Steepest-descent moment method for three-dimensional magnetohydrodynamic equilibria. *The Physics of fluids* **26** (12), 3553–3568.
- HUANG, X., MORITA, S., OISHI, T., GOTO, M. & ZHANG, H. 2015 Evaluation of zeff profile in low-density and high-ti discharges with carbon pellet injection based on space-resolved euv spectroscopy in lhd. *Plasma and Fusion Research* **10**, 3402036–3402036.
- IDA, K., YOSHINUMA, M., KOBAYASHI, T., FUJIWARA, Y., CHEN, J., MURAKAMI, I., KISAKI, M., OSAKABE, M. & LHD EXPERIMENT GROUP 2019 Verification of carbon density profile measurements with charge exchange spectroscopy using hydrogen and deuterium neutral beams. *Plasma and Fusion Research* **14**, 1402079–1402079.
- ISHIKAWA, M., TAKECHI, M., SHINOHARA, K., CHENG, C.-Z., MATSUNAGA, G., KUSAMA, Y., FUKUYAMA, A., NISHITANI, T., MORIOKA, A., SASAO, M., BABA, M. & LHD EXPERIMENT GROUP 2007 Confinement degradation and transport of energetic ions due to alfvén eigenmodes in jt-60u weak shear plasmas. *Nuclear Fusion* **47** (8), 849.
- ISOBE, M., OGAWA, K., NISHITANI, T., MIYAKE, H., KOBUCHI, T., PU, N., KAWASE, H., TAKADA, E., TANAKA, T., LI, S., YOSHIHASHI, S., URITANI, A., JO, J., MURAKAMI, S., OSAKABE, M. & LHD EXPERIMENT GROUP 2018 Neutron diagnostics in the large helical device. *IEEE Transactions on Plasma Science* **46**, 2050–2058.
- ISOBE, M., SASAO, M., OKAMURA, S., KONDO, T., MURAKAMI, S., MINAMI, T., KADO, S., IDA, K., SHIMIZU, A., OSAKABE, M., YOSHINUMA, Y., TANAKA, K., TAKAHASHI, C., NISHIMURA, S., MATSUOKA, K. & CHS GROUP 2001 Experimental investigation of the ripple induced losses of perpendicularly injected beam ions in the low aspect ratio helical system chs. *Nuclear fusion* **41** (9), 1273.
- KAMIO, S., FUJIWARA, Y., OGAWA, K., ISOBE, M., SEKI, R., NUGA, H., NISHITANI, T., OSAKABE, M. & LHD EXPERIMENT GROUP 2019 Development of npa array using single crystal cvd diamond detectors. *Journal of Instrumentation* **14** (08), C08002.
- KARNEY, C. F. F. 1986 “FOKKER-PLANCK AND QUASILINEAR CODES”. *Computer Physics Reports* **4**, 183–244.
- KAWAHATA, K., PETERSON, B. J., AKIYAMA, T., ASHIKAWA, N., EMOTO, M., FUNABA, H., HAMADA, Y., IDA, K., INAGAKI, S., IDO, T., ISOBE, M., GOTO, M., MASE, A., MASUZAKI, S., MICHAEL, C., MORISAKI, T., MORITA, S., MUTO, S., NAGAYAMA, Y., NAKAMURA, Y., NAKANISHI, H., SAKAMOTO, R., NARIHARA, K., NISHIURA, M., OHDACHI, S., OKAJIMA, S., OSAKABE, M., SAKAKIBARA, S., SANIN, A., SASAO, M., SATO, K., SHIMIZU, A., SHOJI, M., SUDO, S., TAMURA, N., TANAKA, K., TOI, K., TOKUZAWA, T., VESHCHEV, E. V., VYACHESLAVOV, L. N., YAMADA, I., YOSHINUMA, M. & LHD EXPERIMENT GROUP 2010 Overview of lhd plasma diagnostics. *Fusion Science and Technology* **58** (1), 331–344.
- LUO, Y., HEIDBRINK, W. W. & BURRELL, K. H. 2004 Design of a d-alpha beam-ion profile diagnostic. *Review of scientific instruments* **75** (10), 3468–3470.
- MEDLEY, S. S. & ROQUEMORE, A. L. 1998 Construction and operation of parallel electric and magnetic field spectrometers for mass/energy resolved multi-ion charge exchange

- diagnostics on the tokamak fusion test reactor. *Review of scientific instruments* **69** (7), 2651–2662.
- MIKKELSEN, D. R. 1989 Approximation for non-resonant beam target fusion reactivities. *Nuclear Fusion* **29** (7), 1113.
- MURAKAMI, S., NAKAJIMA, N. & OKAMOTO, M. 1995 Finite beta effect on the icrf abd nbi heating in the large helical device. *Transactions of Fusion Technology* **27**, 256–259.
- MURAKAMI, S., WAKASA, A., MAASSBERG, H., BEIDLER, C. D., YAMADA, H., WATANABE, K. Y. & LHD EXPERIMENTAL GROUP 2002 Neoclassical transport optimization of lhd. *Nuclear fusion* **42** (11), L19.
- NARIHARA, K., YAMADA, I., HAYASHI, H. & YAMAUCHI, K. 2001 Design and performance of the thomson scattering diagnostic on lhd. *Review of Scientific Instruments* **72** (1), 1122–1125.
- NUGA, H., SEKI, R., OGAWA, K., KAMIO, S., FUJIWARA, Y., OSAKABE, M., ISOBE, M., NISHITANI, T., YOKOYAMA, M. & LHD EXPERIMENT GROUP 2019 Analysis of energetic particle confinement in lhd using neutron measurement and simulation codes. *Plasma and Fusion Research* **14**, 3402075–3402075.
- OGAWA, K., ISOBE, M., NISHITANI, T., MURAKAMI, S., SEKI, R., NUGA, H., KAMIO, S., FUJIWARA, Y., YAMAGUCHI, H., SAITO, Y., MAETA, S., OSAKABE, M. & LHD EXPERIMENT GROUP 2019 Energetic ion confinement studies using comprehensive neutron diagnostics in the large helical device. *Nuclear Fusion* **59** (7), 076017.
- OGAWA, K., ISOBE, M., TOI, K., WATANABE, F., SPONG, D. A., SHIMIZU, A., OSAKABE, M., OHDAKI, S., SAKAKIBARA, S. & LHD EXPERIMENT GROUP 2010 Observation of energetic-ion losses induced by various mhd instabilities in the large helical device (lhd). *Nuclear Fusion* **50** (8), 084005.
- OSAKABE, M., ISOBE, M., TANAKA, M., MOTOJIMA, G., TSUMORI, K., YOKOYAMA, M., MORISAKI, T. & TAKEIRI, Y. 2018 Preparation and commissioning for the lhd deuterium experiment. *IEEE Transactions on Plasma Science* **46**, 2324–2331.
- OSAKABE, M., TAKEIRI, Y., MORISAKI, T., MOTOJIMA, G., OGAWA, K., ISOBE, M., TANAKA, M., MURAKAMI, S., SHIMIZU, A., NAGAOKA, K., TAKAHASHI, H., NAGASAKI, K., TAKAHASHI, H., FUJITA, T., OYA, Y., SAKAMOTO, M., UEDA, Y., AKIYAMA, T., KASAHARA, H., SAKAKIBARA, S., SAKAMOTO, R., TOKITANI, M., YAMADA, H., YOKOYAMA, M., YOSHIMURA, Y. & LHD EXPERIMENT GROUP 2017 Current status of large helical device and its prospect for deuterium experiment. *Fusion Science and Technology* **72** (3), 199–210.
- OSAKABE, M., YAMAMOTO, S., TOI, K., TAKEIRI, Y., SAKAKIBARA, S., NAGAOKA, K., TANAKA, K., NARIHARA, K. & LHD EXPERIMENTAL GROUP 2006 Experimental observations of enhanced radial transport of energetic particles with alfvén eigenmode on the lhd. *Nuclear fusion* **46** (10), S911.
- RUSKOV, E., BELL, M., BUDNY, R. V., MCCUNE, D. C., MEDLEY, S. S., REDI, M. H., SCOTT, S., SYNAKOWSKI, E. J., VON GOELER, S., WHITE, R. B. & ZWEBEN, S. J. 1999 Anomalous beam-ion loss in tfr reversed magnetic shear plasmas. *Phys. Rev. Lett.* **82**, 924–927.
- SATO, M., MURAKAMI, S., FUKUYAMA, A., NAKAMURA, Y., WATANABE, K. Y., TODA, S., YOKOYAMA, M., FUNABA, H., YAMADA, H. & NAKAJIMA, N. 2008 Implementation of nbi heating module fit3d to hierarchy-integrated simulation code task3d. In *Proc. 18th Int. Toki Conf.*
- SEKI, R., OGAWA, K., ISOBE, M., YOKOYAMA, M., MURAKAMI, S., NUGA, H., KAMIO, S., FUJIWARA, Y., OSAKABE, M. & LHD EXPERIMENT GROUP 2019 Evaluation of neutron emission rate with fit3d-dd code in large helical device. *Plasma and Fusion Research* **14**, 3402126–3402126.
- STRACHAN, J. D., COLESTOCK, P. L., DAVIS, S. L., EAMES, D., EFTHIMION, P. C., EUBANK, H. P., GOLDSTON, R. J., GRISHAM, L. R., HAWRYLUK, R. J., HOSEA, J. C., HOVEY, J., JASSBY, D. L., JOHNSON, D. W., MIRIN, A. A., SCHILLING, G., STOOKSBERRY, R., STEWART, L. D. & TOWNER, H. H. 1981 Fusion neutron production during deuterium neutral-beam injection into the plt tokamak. *Nuclear Fusion* **21** (1), 67.
- TAKEIRI, Y. 2018a The large helical device: Entering deuterium experiment phase toward steady-state helical fusion reactor based on achievements in hydrogen experiment phase. *IEEE Transactions on Plasma Science* **46**, 2348–2353.

- TAKEIRI, Y. 2018*b* Prospect toward steady-state helical fusion reactor based on progress of lhd project entering the deuterium experiment phase. *IEEE Transactions on Plasma Science* **46** (5), 1141–1148.
- TAKEIRI, Y., KANEKO, O., TSUMORI, K., OSAKABE, M., IKEDA, K., NAGAOKA, K., NAKANO, H., ASANO, E., KONDO, T., SATO, M., SHIBUYA, M., KOMADA, S. & LHD EXPERIMENT GROUP 2010 High performance of neutral beam injectors for extension of lhd operational regime. *Fusion Science and Technology* **58** (1), 482–488.
- TOBITA, K., TANI, K., NISHITANI, T., NAGASHIMA, K. & KUSAMA, Y. 1994 Fast ion losses due to toroidal field ripple in jt-60u. *Nuclear fusion* **34** (8), 1097.
- VINCENZI, P., BOLZONELLA, T., MURAKAMI, S., OSAKABE, M., SEKI, R. & YOKOYAMA, M. 2016 Upgrades and application of fit3d nbi–plasma interaction code in view of lhd deuterium campaigns. *Plasma Physics and Controlled Fusion* **58** (12), 125008.
- YAMADA, I., NARIHARA, K., FUNABA, H., MINAMI, T., HAYASHI, H., KOHMOTO, T. & LHD EXPERIMENT GROUP 2010 Recent progress of the lhd thomson scattering system. *Fusion Science and Technology* **58** (1), 345–351.
- YOKOYAMA, M., SEKI, R., SUZUKI, C., SATO, M., EMOTO, M., MURAKAMI, S., OSAKABE, M., TSUJIMURA, T. I., YOSHIMURA, Y., IDO, T., OGAWA, K., SATAKE, S., SUZUKI, Y., GOTO, T., IDA, K., PABLANT, N., GATES, D., WARMER, F., VINCENZI, P. & NUMERICAL SIMULATION REACTOR RESEARCH PROJECT AND LHD EXPERIMENT GROUP 2017 Extended capability of the integrated transport analysis suite, task3d-a, for lhd experiment. *Nuclear Fusion* **57** (12), 126016.
- ZHANG, Y. P., ISOBE, M., LIU, Y., YUAN, G. L., YANG, J. W., SONG, X. Y., SONG, X. M., CAO, J. Y., LEI, G. J., WEI, H. L., LI, Y. G., SHI, Z. B., LI, X., YAN, L. W., YANG, Q. W., DUAN, X. R. & HL-2A TEAM 2012 Measurements of the fast ion slowing-down times in the hl-2a tokamak and comparison to classical theory. *Physics of Plasmas* **19** (11), 112504.
- ZHOU, H., MORITA, S., GOTO, M., DONG, C., YANAGIBAYASHI, J. & HASUO, M. 2010 Z_{eff} measurement using extreme ultraviolet bremsstrahlung emission in lhd. *Review of Scientific Instruments* **81** (10), 10D706.