§22. Theory of Anomalous Transport in H-Mode Plasmas

Itoh, S.-I. (Kyushu Univ.)
 Itoh, K.
 Fukuyama, A. (Okayama Univ.)
 Yagi, M. (JAERI)

The understanding of the transport in H-mode ultimately requires that of the L-mode plasma simultaneously. We apply the method of self-sustaining turbulence to the tokamak plasma with the inhomogeneous E_r . The anomalous transport coefficient is obtained in an explicit form of the parameters which characterize the plasma profile. Improvement of the thermal conductivity in the H-mode plasma is explained. Change in the fluctuations is also analyzed.

The basic equation which describe the self-sustaining turbulence in the H-mode plasma was derived [1]. By solving the eigen value equation in terms of the transport coefficient, we have the anomalous transport coefficient in the presence of the inhomogeneous radial electric field is obtained

$$\chi = \frac{f(s)^{-1} \alpha^{3/2} (\lambda/\tilde{\chi}) (\tilde{\chi}/\tilde{\mu})^{1/2}}{1 + G_1 \omega_{E1}^2} \quad (1)$$

where

$$G_1 = \left[\frac{9}{8} \left(1 + \frac{4}{3\alpha f} \right) + \frac{25s^2}{4(1+2\alpha-2s)} \sqrt{\frac{2+C}{1+C}} \left(1 + \frac{4}{3\alpha f} \right) \left(1 + \frac{4}{5\alpha f} \right) \right] \frac{\alpha}{f^2}$$

and

$$\omega_{E1} = \tau_{A0} (dE_r/d\tilde{r}) (srB)^{-1}.$$

(Length and time are normalized to a and $\tau_{Ap} \equiv a\sqrt{\mu_0 m_i n_i}/B_p$.)

The numerator of Eq. (1) is the transport coefficient in the L-mode, χ_L [2]. Equation (1) quantifies the effect of E_r on the thermal conductivity, unifying the L- and H-mode plasmas. It is emphasized that the coefficients are given explicitly in terms of the equilibrium quantities. This is because the self-sustained turbulence is solved by our theoretical formalism. The suppression of the transport is prominent when $\omega_{E1} \sim 1/\sqrt{G_1}$. A noticeable reduction of χ occurs when ω_{E1} approaches unity. The coefficient f denotes the influence of the magnetic shear.

The theory also predicts the change in the fluctuations. The estimate of k_θ is given as

$$k_\theta^2 = \delta^{-2} \alpha^{-1} \left[1 + G_1 \omega_{E1}^2 \right], \quad (2)$$

where δ is the collisionless skin depth. Fluctuation level is also reduced.

$$e\tilde{\phi}/T \sim (\chi_L eB/T) \left[1 + G_1 \omega_{E1}^2 \right]^{-1}. \quad (3)$$

These results present the unified description on the transport coefficient and fluctuations in the H-mode as well as in the L-mode plasmas.

References

- [1] K. Itoh et al.: "Effect of Flow shear on Current-Diffusive Ballooning Mode", in this issue.
- [2] K. Itoh, et al.: Plasma Phys. and Control. Fusion **35** (1993) 543.
- [3] S.-I. Itoh et al.: Phys. Rev. Lett. **72** (1994) 1200.