

§11. H-mode-Like Discharges and ELM-Like Activities under the Presence of $\iota/2\pi=1$ at Ergodic Layer in LHD

Morita, S., Morisaki, T., Tanaka, K., Masuzaki, S., Goto, M., Sakakibara, S., Watanabe, K.Y.

Shifting the R_{ax} outwardly, the profile of rotational transform becomes flat and the edge rotational transform at $\rho=1$, $\iota/2\pi(a)$, becomes small with decreasing the edge magnetic shear (see Fig.1). Then, the values of $\iota/2\pi(a)$ are below unity at $R_{ax} \geq 3.90\text{m}$. The H-mode-like discharge with ELM-like activity has been observed in LHD at outwardly shifted configuration where $\iota/2\pi=1$ exists inside the edge ergodic layer.

A typical waveform of the H-mode-like discharge is shown in Fig.2. The H-mode-like transition is triggered by changing the NBI power input at $t=1.25\text{s}$. After turning off the beam line, the $H\alpha$ emission quickly drops and the density gradually rises, showing a clear turning point. ELM-like bursts appear in the $H\alpha$ signal. Similar bursts are also observed in an electrostatic probe on the divertor plate (I_{is}) and a magnetic probe (db/dt). Reduction of the magnetic fluctuation is also seen after the H-mode-like transition. No clear mode numbers are observed on the ELM-like bursts from the magnetic probe measurement. This H-mode-like feature, however, disappears after turning off the second NBI at $t=2.1\text{s}$. It strongly suggests that a relatively narrow power window exists for keeping the H-mode-like phase.

Pressure profiles in the ergodic layer measured with YAG laser Thomson diagnostics are plotted in Fig.3. The data are obtained at $t=1.225\text{s}$ before transition (open circles) and $t=1.625\text{s}$ after transition (solid circles) in the same discharge shown in Fig.2. The position of $\iota/2\pi=1$ surface is indicated by a vertical dashed line and the LCFS position is $R=4.560\text{m}$ shown by a vertical solid line. The edge pressure rapidly increases near $\iota/2\pi=1$ position and it is seen that a steeper pressure gradient is formed after the transition, although the edge T_e decreases after the transition because of density increase.

Temporal behaviors of three chord-integrated densities from the edge region are traced in Fig.4(a)-(c). The ergodic layer becomes thick at the inboard side, then, the signals of (a) and (b) show the information only from the ergodic layer. A relatively high density exists even in the middle of the ergodic layer of $R=3.399\text{m}$ where the L_c is roughly 100m . A high density is not formed at a region of $L_c < 10\text{m}$. Precise measurement was also done to investigate the relation between the $\iota/2\pi=1$ position and the density bursts position using a multichannel CO_2 interferometer with a spatial resolution of 1cm . It is seen that the amplitude of the density bursts become the largest at $\iota/2\pi=1$ position.

The edge rotational transform in vacuum is used at present. Taking into account the plasma pressure, the edge rotational transform profile seems to be modified, based on a modification of chaotic magnetic field structure

of the ergodic layer. A detailed discussion has to be made for the modification.




Fig. 1 Rotational transform as a parameter of R_{ax} .




Fig. 2 H-mode-like discharge with ELM-like bursts.




Fig. 3 Edge pressure profile before and after transition.




Fig. 4 Edge density behaviors during ELM-like bursts.

References

- 1) Morita, S. et al., PPCF **48** (2006) A269.