



INTERNATIONAL ATOMIC ENERGY AGENCY

**FIFTEENTH INTERNATIONAL CONFERENCE ON PLASMA PHYSICS
AND CONTROLLED NUCLEAR FUSION RESEARCH**

Seville, Spain, 26 September – 1 October 1994

IAEA-CN-60/A-2-IV-2

NATIONAL INSTITUTE FOR FUSION SCIENCE
**Control of Radial Electric Field in
Torus Plasma**

K. Ida, H. Idei, H. Sanuki, K. Itoh, J. Xu, S. Hidekuma, K. Kondo, A. Sahara, H. Zushi, S.-I. Itoh, A. Fukuyama, K. Adati, R. Akiyama, S. Bessho, A. Ejiri, A. Fujisawa, M. Fujiwara, Y. Hamada, S. Hirokura, H. Iguchi, O. Kaneko, K. Kawahata, Y. Kawasumi, M. Kojima, S. Kubo, H. Kuramoto, A. Lazaros, R. Liang, K. Matsuoka, T. Minami, T. Mizuuchi, T. Morisaki, S. Morita, K. Nagasaki, K. Narihara, K. Nishimura, A. Nishizawa, T. Obiki, H. Okada, S. Okamura, T. Ozaki, S. Sakakibara, H. Sakakita, A. Sagara, F. Sano, M. Sasao, K. Sato, K.N. Sato, T. Saeki, S. Sudo, C. Takahashi, K. Takaka, K. Tsumori, H. Yamada, I. Yamada, Y. Takita, T. Tuzuki, K. Toi and T. Watari
(Received - Aug. 26, 1994)

NIFS-304

Sep. 1994

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Nagoya 464-01, Japan.

**RESEARCH REPORT
NIFS Series**

This is a preprint of a paper intended for presentation at a scientific meeting. Because of the provisional nature of its contents and since changes of substance or detail may result by the publication, the preprint is made available on the understanding that it will remain cited in the literature or in any way be reproduced in its present form. The views expressed and the statements made remain the responsibility of the named author(s). The views do not necessarily reflect those of the government of the designating Member State(s) or of the designating organization(s). In particular, neither the IAEA nor any other organization of which the author(s) are a member(s) are held responsible for any material reproduced in this preprint.

NAGOYA, JAPAN



CONTROL OF RADIAL ELECTRIC FIELD IN TORUS PLASMA

K.IDA, H.IDEI, H.SANUKI, K.ITOH, J.XU, S.HIDEKUMA, K.KONDO¹, A.SAHARA¹, H.ZUSHI¹, S.-I.ITOH², A.FUKUYAMA³, K.ADATI, R.AKIYAMA, S.BESSHO¹, A.EJIRI, A.FUJISAWA, M.FUJIWARA, Y.HAMADA, S.HIROKURA, H.IGUCHI, O.KANEKO, K.KAWAHATA, Y.KAWASUMI, M.KOJIMA, S.KUBO, H.KURAMOTO, A.LAZAROS⁴, R.LIANG, K.MATSUOKA, T.MINAMI, T.MIZUUCHI¹, T.MORISAKI, S.MORITA, K.NAGASAKI¹, K.NARIHARA, K.NISHIMURA, A.NISHIZAWA, T.OBIKI¹, H.OKADA¹, S.OKAMURA, T.OZAKI, S.SAKAKIBARA, H.SAKAKITA, A.SAGARA, F.SANO¹, M.SASAO, K.SATO, K.N.SATO, T.SEKI, S.SUDO, C.TAKAHASHI, K.TANAKA, K.TSUMORI, H.YAMADA, I.YAMADA, Y.TAKITA, T.TUZUKI, K.TOI, T.WATARI

National Institute for Fusion Science 464-01 Nagoya Japan

¹ Plasma Physics Laboratory, Kyoto University, Uji, Kyoto, 611 Japan

² Research Institute for Applied Mechanics, Kyusu University, Kasuga, 816 Japan

³ Faculty of engineering, Okayama University, Okayama, 700 Japan

⁴ European Community Science and Technology Program

CONTROL OF RADIAL ELECTRIC FIELD IN TORUS PLASMA

Abstract

The radial electric field is controlled by changing the direction of neutral beam from co to counter to plasma current in tokamak, while it is controlled by the 2nd harmonic ECH and NBI and pellet injection in heliotron/torsatron.

Keywords : Radial electric field, torus plasma

1. INTRODUCTION

Radial electric field E_r is considered to be one of the key parameter to improve plasma confinement. In order to control E_r profiles, the electron cyclotron resonance heating (ECH) and/or neutral beam injection (NBI) are applied in tokamak and heliotron/torsatron plasma. Detailed radial profiles of electric field are obtained from the radial profiles of toroidal and poloidal rotation velocity measured with multi-channel charge exchange spectroscopy (CXs).

2. CONTROL OF RADIAL ELECTRIC FIELD BY NEUTRAL BEAM INJECTION(NBI) IN JIPP TII-U AND CHS

In the tokamak, the toroidal rotation velocity profile can be controlled by switching the direction of momentum input of high energy neutral beam from parallel (co-injection) to anti-parallel (counter-injection) to the direction of plasma current. Figure 1 shows the radial electric field profiles normalized by central ion temperature for parallel injection and near perpendicular injection (9 degree) in JIPP TIIU tokamak. The radial electric field is negative for perpendicular NB injected plasma. It becomes positive near the plasma center for co-injection, while it becomes large negative value for counter-injected NB plasma. This result illustrates the offset part of rotation (spontaneous rotation) and the combination of the external torque. Since the offset toroidal rotation velocity is in the counter-direction (negative electric field), the absolute value of electric field is much larger in counter-injected plasma than that in co-injected plasma[1]. Density peaking associated with this negative electric field indicates the improvement of particle transport in tokamak.




Fig.1 Radial electric field profile in JIPP T-IIU tokamak, where a is plasma minor radius




Fig.2 Radial electric field profile in CHS heliotron/torsatron, where a is plasma minor radius.

On the other hand in CHS heliotron/torsatron, even parallel NBI can not drive toroidal plasma rotation large enough to change radial electric field, since parallel viscosity is dominant in the configuration ($R_{ax} = 97.4\text{cm}$: magnetic field ripple of 2-3% even at the magnetic axis)[2]. The radial electric field in high density plasma is negative regardless of the direction of neutral beam in heliotron/torsatron as shown in Figure 2. The radial electric field depends on other bipolar fluxes such

as ion orbit loss, charge-exchange loss, neoclassical and anomalous ion/electron fluxes; it does not depend on the NB injection direction dominantly. When the magnetic axis is shifted toward the direction in which more fast ion loss is expected (inward for ctr-NBI and outward for co-NBI), more negative electric field is observed. When the electron density is decreased below $1 \times 10^{13} \text{cm}^{-3}$, the negative radial electric field becomes smaller as expected by neoclassical theory, but no positive radial electric field is observed in NB heated plasma in CHS.

3. CONTROL OF RADIAL ELECTRIC FIELD BY ELECTRON CYCLOTRON RESONANCE HEATING (ECH) IN CHS

The radial electric field is more easily controlled by use of bipolar fluxes due to ECH than the momentum input of NBI in CHS because of magnetic field ripple. The electron cyclotron resonance heating (ECH) has been considered to enhance positive radial electric field, because it heats the electron and enhances electron ripple loss. In order to produce positive electric field, 53GHz ECH is injected to the NBI plasma with the helical field of 0.92 and 1.9T in CHS. The experiment is restricted only for the NB target plasma, because charge-exchange measurement needs NBI. The positive radial electric field is observed only when the off-axis second harmonic ECH is applied to the low density ($n_e < 1 \times 10^{13} \text{cm}^{-3}$) NB heated plasma[3]. As seen in Fig.3 there is a critical ECH power for the radial electric field to jump to positive values [electron root]. This jump of radial electric field is more visible near the plasma edge and is considered to be due to the ECH enhanced electron loss. No positive radial electric field is observed even if the plasma with similar electron density and temperature is sustained only by NBI. The perpendicular acceleration of ripple trapped electron is one of the candidate of this electron loss.




Fig.3 Radial electric field at each averaged minor radius as a function of ECH power in CHS.




Fig.4 Radial electric field profiles for ECH and for ECH plus ctr-NBI in CHS

The ion dynamics would be also important in determining the total radial electric field. As shown in Figure 4, the positive radial electric field produced by ECH disappears when the neutral beam is injected in counter direction. No improvement of energy transport by the positive electric field is observed at the transition to the electron-root. This is because the radial electric field shear is not large enough to suppress fluctuations. However, if the viscosity for plasma rotation decreases by the reduction of edge neutral density, or the focusing of the ECH power becomes better, the larger radial electric field

shear will be produced and the improvement of energy confinement also will be expected.

4. CONTROL OF RADIAL ELECTRIC FIELD BY PELLET INJECTION IN HELIOTRON-E

Radial electric field profiles are measured before and after the pellet injection for NBI heated plasmas in Heliotron-E. The E_r profile before the pellet injection is similar to that for NBI heated plasma in CHS (more negative toward the plasma edge). However, as shown in Fig.5, E_r becomes more negative and has a peak at the half of plasma minor radius after the pellet injection, where the electron density is more peaked at the plasma center. This experiment suggests that the E_r can be controlled through the density profile with pellet injection.




Fig. 5. Radial electric field profiles before and after the pellet injection in heliotron-E.

5. SUMMARY

In conclusion, the radial electric field is controlled by changing the direction of neutral beam from co to counter to plasma current in tokamak, while it is controlled by the injection of 2nd harmonic ECH and NBI (either co or counter direction) and pellet injection in heliotron/torsatron.

REFERENCE

- [1] IDA, K., et al., Nucl. Fusion **31** (1991) 943.
- [2] IDA, K., YAMADA, H., IGUCHI, H.i, et al., Phys. Rev. Lett. **67** (1991) 58.
- [3] IDEI, H., IDA, K., SANUKI, H., et al., Phys. Rev. Lett. **71** (1993) 2220.

Recent Issues of NIFS Series

- NIFS-264 K. Masai,
Nonequilibria in Thermal Emission from Supernova Remnants;
Dec. 1993
- NIFS-265 K. Masai, K. Nomoto,
*X-Ray Enhancement of SN 1987A Due to Interaction with its Ring-like
Nebula;* Dec. 1993
- NIFS-266 J. Uramoto
*A Research of Possibility for Negative Muon Production by a Low
Energy Electron Beam Accompanying Ion Beam;* Dec. 1993
- NIFS-267 H. Iguchi, K. Ida, H. Yamada, K. Itoh, S.-I. Itoh, K. Matsuoka,
S. Okamura, H. Sanuki, I. Yamada, H. Takenaga, K. Uchino, K. Muraoka,
*The Effect of Magnetic Field Configuration on Particle Pinch Velocity in
Compact Helical System (CHS);* Jan. 1994
- NIFS-268 T. Shikama, C. Namba, M. Kosuda, Y. Maeda,
*Development of High Time-Resolution Laser Flash Equipment for
Thermal Diffusivity Measurements Using Miniature-Size Specimens;*
Jan. 1994
- NIFS-269 T. Hayashi, T. Sato, P. Merkel, J. Nührenberg, U. Schwenn,
*Formation and 'Self-Healing' of Magnetic Islands in Finite- β Helias
Equilibria;* Jan. 1994
- NIFS-270 S. Murakami, M. Okamoto, N. Nakajima, T. Mutoh,
*Efficiencies of the ICRF Minority Heating in the CHS and LHD
Plasmas;* Jan. 1994
- NIFS-271 Y. Nejoh, H. Sanuki,
*Large Amplitude Langmuir and Ion-Acoustic Waves in a Relativistic
Two-Fluid Plasma;* Feb. 1994
- NIFS-272 A. Fujisawa, H. Iguchi, A. Taniike, M. Sasao, Y. Hamada,
A 6MeV Heavy Ion Beam Probe for the Large Helical Device;
Feb. 1994
- NIFS-273 Y. Hamada, A. Nishizawa, Y. Kawasumi, K. Narihara, K. Sato, T. Seki, K. Toi,
H. Iguchi, A. Fujisawa, K. Adachi, A. Ejiri, S. Hidekuma, S. Hirokura,
K. Ida, J. Koong, K. Kawahata, M. Kojima, R. Kumazawa, H. Kuramoto,
R. Liang, H. Sakakita, M. Sasao, K. N. Sato, T. Tsuzuki, J. Xu, I. Yamada,
T. Watari, I. Negi,
*Measurement of Profiles of the Space Potential in JIPP T-IIU Tokamak
Plasmas by Slow Poloidal and Fast Toroidal Sweeps of a Heavy Ion
Beam;* Feb. 1994

- NIFS-274 M. Tanaka,
A Mechanism of Collisionless Magnetic Reconnection; Mar. 1994
- NIFS-275 A. Fukuyama, K. Itoh, S.-I. Itoh, M. Yagi and M. Azumi,
Isotope Effect on Confinement in DT Plasmas; Mar. 1994
- NIFS-276 R.V. Reddy, K. Watanabe, T. Sato and T.H. Watanabe,
Impulsive Alfvén Coupling between the Magnetosphere and Ionosphere; Apr. 1994
- NIFS-277 J. Uramoto,
A Possibility of π^- Meson Production by a Low Energy Electron Bunch and Positive Ion Bunch; Apr. 1994
- NIFS-278 K. Itoh, S.-I. Itoh, A. Fukuyama, M. Yagi and M. Azumi,
Self-sustained Turbulence and L-mode Confinement in Toroidal Plasmas II; Apr. 1994
- NIFS-279 K. Yamazaki and K.Y. Watanabe,
New Modular Heliotron System Compatible with Closed Helical Divertor and Good Plasma Confinement; Apr. 1994
- NIFS-280 S. Okamura, K. Matsuoka, K. Nishimura, K. Tsumori, R. Akiyama, S. Sakakibara, H. Yamada, S. Morita, T. Morisaki, N. Nakajima, K. Tanaka, J. Xu, K. Ida, H. Iguchi, A. Lazaros, T. Ozaki, H. Arimoto, A. Ejiri, M. Fujiwara, H. Idei, O. Kaneko, K. Kawahata, T. Kawamoto, A. Komori, S. Kubo, O. Motojima, V.D. Pustovitov, C. Takahashi, K. Toi and I. Yamada,
High-Beta Discharges with Neutral Beam Injection in CHS; Apr. 1994
- NIFS-281 K. Kamada, H. Kinoshita and H. Takahashi,
Anomalous Heat Evolution of Deuteron Implanted Al on Electron Bombardment ; May 1994
- NIFS-282 H. Takamaru, T. Sato, K. Watanabe and R. Horiuchi,
Super Ion Acoustic Double Layer; May 1994
- NIFS-283 O. Mitarai and S. Sudo
Ignition Characteristics in D-T Helical Reactors; June 1994
- NIFS-284 R. Horiuchi and T. Sato,
Particle Simulation Study of Driven Magnetic Reconnection in a Collisionless Plasma; June 1994
- NIFS-285 K.Y. Watanabe, N. Nakajima, M. Okamoto, K. Yamazaki, Y. Nakamura, M. Wakatani,
Effect of Collisionality and Radial Electric Field on Bootstrap Current in LHD (Large Helical Device); June 1994

- NIFS-286 H. Sanuki, K. Itoh, J. Todoroki, K. Ida, H. Idei, H. Iguchi and H. Yamada,
*Theoretical and Experimental Studies on Electric Field and Confinement
in Helical Systems*; June 1994
- NIFS-287 K. Itoh and S.-I. Itoh,
Influence of the Wall Material on the H-mode Performance; June 1994
- NIFS-288 K. Itoh, A. Fukuyama, S.-I. Itoh, M. Yagi and M. Azumi
Self-Sustained Magnetic Braiding in Toroidal Plasmas: July 1994
- NIFS-289 Y. Nejoh,
*Relativistic Effects on Large Amplitude Nonlinear Langmuir Waves
in a Two-Fluid Plasma*; July 1994
- NIFS-290 N. Ohyabu, A. Komori, K. Akaishi, N. Inoue, Y. Kubota, A.I. Livshit,
N. Noda, A. Sagara, H. Suzuki, T. Watanabe, O. Motojima, M. Fujiwara,
A. Iiyoshi,
Innovative Divertor Concepts for LHD; July 1994
- NIFS-291 H. Idei, K. Ida, H. Sanuki, S. Kubo, H. Yamada, H. Iguchi, S. Morita,
S. Okamura, R. Akiyama, H. Arimoto, K. Matsuoka, K. Nishimura,
K. Ohkubo, C. Takahashi, Y. Takita, K. Toi, K. Tsumori and I. Yamada,
*Formation of Positive Radial Electric Field by Electron Cyclotron
Heating in Compact Helical System*; July 1994
- NIFS-292 N. Noda, A. Sagara, H. Yamada, Y. Kubota, N. Inoue, K. Akaishi, O. Motojima,
K. Iwamoto, M. Hashiba, I. Fujita, T. Hino, T. Yamashina, K. Okazaki,
J. Rice, M. Yamage, H. Toyoda and H. Sugai,
Boronization Study for Application to Large Helical Device; July 1994
- NIFS-293 Y. Ueda, T. Tanabe, V. Philipps, L. Könen, A. Pospieszczyk, U. Samm,
B. Schweer, B. Unterberg, M. Wada, N. Hawkes and N. Noda,
*Effects of Impurities Released from High Z Test Limiter on Plasma
Performance in TEXTOR*; July. 1994
- NIFS-294 K. Akaishi, Y. Kubota, K. Ezaki and O. Motojima,
*Experimental Study on Scaling Law of Outgassing Rate with A Pumping
Parameter*, Aug. 1994
- NIFS-295 S. Bazdenkov, T. Sato, R. Horiuchi, K. Watanabe
*Magnetic Mirror Effect as a Trigger of Collisionless Magnetic
Reconnection*, Aug. 1994
- NIFS-296 K. Itoh, M. Yagi, S.-I. Itoh, A. Fukuyama, H. Sanuki, M. Azumi
Anomalous Transport Theory for Toroidal Helical Plasmas,
Aug. 1994 (IAEA-CN-60/D-III-3)

- NIFS-297 J. Yamamoto, O. Motojima, T. Mito, K. Takahata, N. Yanagi, S. Yamada, H. Chikaraishi, S. Imagawa, A. Iwamoto, H. Kaneko, A. Nishimura, S. Satoh, T. Satow, H. Tamura, S. Yamaguchi, K. Yamazaki, M. Fujiwara, A. Iiyoshi and LHD group,
New Evaluation Method of Superconductor Characteristics for Realizing the Large Helical Device; Aug. 1994 (IAEA-CN-60/F-P-3)
- NIFS-298 A. Komori, N. Ohyabu, T. Watanabe, H. Suzuki, A. Sagara, N. Noda, K. Akaishi, N. Inoue, Y. Kubota, O. Motojima, M. Fujiwara and A. Iiyoshi,
Local Island Divertor Concept for LHD; Aug. 1994 (IAEA-CN-60/F-P-4)
- NIFS-299 K. Toi, T. Morisaki, S. Sakakibara, A. Ejiri, H. Yamada, S. Morita, K. Tanaka, N. Nakajima, S. Okamura, H. Iguchi, K. Ida, K. Tsumori, S. Ohdachi, K. Nishimura, K. Matsuoka, J. Xu, I. Yamada, T. Minami, K. Narihara, R. Akiyama, A. Ando, H. Arimoto, A. Fujisawa, M. Fujiwara, H. Idei, O. Kaneko, K. Kawahata, A. Komori, S. Kubo, R. Kumazawa, T. Ozaki, A. Sagara, C. Takahashi, Y. Takita and T. Watari
Impact of Rotational-Transform Profile Control on Plasma Confinement and Stability in CHS; Aug. 1994 (IAEA-CN-60/A6/C-P-3)
- NIFS-300 H. Sugama and W. Horton,
Dynamical Model of Pressure-Gradient-Driven Turbulence and Shear Flow Generation in L-H Transition; Aug. 1994 (IAEA/CN-60/D-P-I-11)
- NIFS-301 Y. Hamada, A. Nishizawa, Y. Kawasumi, K.N. Sato, H. Sakakita, R. Liang, K. Kawahata, A. Ejiri, K. Narihara, K. Sato, T. Seki, K. Toi, K. Itoh, H. Iguchi, A. Fujisawa, K. Adachi, S. Hidekuma, S. Hirokura, K. Ida, M. Kojima, J. Koog, R. Kumazawa, H. Kuramoto, T. Minami, I. Negi, S. Ohdachi, M. Sasao, T. Tsuzuki, J. Xu, I. Yamada, T. Watari,
Study of Turbulence and Plasma Potential in JIPP T-IIU Tokamak; Aug. 1994 (IAEA/CN-60/A-2-III-5)
- NIFS-302 K. Nishimura, R. Kumazawa, T. Mutoh, T. Watari, T. Seki, A. Ando, S. Masuda, F. Shinpo, S. Murakami, S. Okamura, H. Yamada, K. Matsuoka, S. Morita, T. Ozaki, K. Ida, H. Iguchi, I. Yamada, A. Ejiri, H. Idei, S. Muto, K. Tanaka, J. Xu, R. Akiyama, H. Arimoto, M. Isobe, M. Iwase, O. Kaneko, S. Kubo, T. Kawamoto, A. Lazaros, T. Morisaki, S. Sakakibara, Y. Takita, C. Takahashi and K. Tsumori,
ICRF Heating in CHS; Sep. 1994 (IAEA-CN-60/A-6-I-4)
- NIFS-303 S. Okamura, K. Matsuoka, K. Nishimura, K. Tsumori, R. Akiyama, S. Sakakibara, H. Yamada, S. Morita, T. Morisaki, N. Nakajima, K. Tanaka, J. Xu, K. Ida, H. Iguchi, A. Lazaros, T. Ozaki, H. Arimoto, A. Ejiri, M. Fujiwara, H. Idei, A. Iiyoshi, O. Kaneko, K. Kawahata, T. Kawamoto, S. Kubo, T. Kuroda, O. Motojima, V.D. Pustovitov, A. Sagara, C. Takahashi, K. Toi and I. Yamada,
High Beta Experiments in CHS; Sep. 1994 (IAEA-CN-60/A-2-IV-3)