

#### §4. Effects of Non-Axisymmetric Magnetic Field on Characteristics of Non-Neutral Plasma

Tomita, Y.,  
Yasaka, Y., Takeno, H. (Kobe Univ.),  
Ishikawa, M., Nakashima, Y., Katanuma, I., Cho, T.  
(Univ. Tsukuba)

In order to investigate the effect of electric field, we considered the axisymmetric hollow equilibrium of the non-neutral plasma, where the densities are uniform. The ions are located at a center region,  $0 \leq r \leq R_i$  and the electrons are cylindrically distributed,  $R_i < R_{e,in} \leq r \leq R_{e,out}$ .

The momentum equation of the non-diamagnetic equilibrium of the  $j$ -th species, where the self magnetic fields are neglected, is expressed as

$$-\frac{m_j V_{j\theta}^2}{r} = q_j (E_r + V_{j\theta} B_z^0). \quad (1)$$

The radial electric field is obtained from the Poisson equation,

$$E_r = \frac{1}{\varepsilon_0 r} \sum_k q_k \int_0^r dr' r' n_k(r'). \quad (2)$$

The momentum equation, Eq.(1), has the quadratic form with respect to the angular frequency  $\omega_j = V_{j\theta}/r$ ,

$$\omega_j^2 + \sigma_j \Omega_j \omega_j + \frac{q_j E_r}{m_j r} = 0. \quad (3)$$

In the case of ions, the equilibrium are obtained by the rigid rotor configuration,

$$\omega_i / \omega_{ci0} = \frac{1}{2} [-1 \pm \sqrt{1 - 2n_{i0} / (\varepsilon_0 B_{z0}^2 / m_i)}], \quad (4)$$

where  $\omega_{ci0}$  is the ion cyclotron frequency to the unperturbed axial magnetic field  $B_{z0}$ . The condition of the real rotation frequency gives the upper limit of the uniform ion density,

$$n_{i0} \leq \frac{\varepsilon_0 B_{z0}^2}{2m_i} = 2.65 \times 10^{15} B_{z0,T}^2 \text{ (m}^{-3}\text{)}, \quad (5)$$

where  $B_{z0,T}$  is the unperturbed axial magnetic field in the unit of Tesla.

For electrons, the rotation frequency is not uniform,

$$\omega_e / \omega_{ce0} =$$

$$\frac{1}{2} [1 \pm \sqrt{1 - \frac{2n_{e0}}{r^2} \frac{\varepsilon_0 B_{z0}^2}{m_e} (R_i^2 - r^2 + R_{e,in}^2)}], \quad (6)$$

which is shown in Fig.1, where  $R_{e,in} / R_i = 2.0$ ,

$R_{e,out} / R_i = 3.0$ , and  $n_{e0} / (\varepsilon_0 B_{z0}^2 / m_e) = 1.0$ , where

$\omega_e^\pm$  corresponds to the sign +/- of the solution in Eq.(6).

This condition satisfies the real condition of the electron angular frequency.

The non-axisymmetric uniform magnetic field  $B_1$  to the vertical direction to the axisymmetric equilibrium configuration changes the radial component of the momentum equation,

$$\omega_j^2 + \sigma_j \Omega_j \omega_j + \frac{q_j}{m_j r} (E_r + V_{jz} B_1 \sin\theta) = 0. \quad (7)$$

One can see that the perturbation changes the effective electric field and deform the axisymmetry. The magnitude of the effect can be estimated by

$$\frac{|V_{jz} B_1|}{E_r} \sim \frac{V_{jz}}{R_i \omega_{cj0}} \frac{B_1}{B_{z0}} \sim \frac{c_s}{R_i \omega_{cj0}} \frac{B_1}{B_{z0}} \sim \frac{B_1}{B_{z0}}, \quad (8)$$

where  $B_{z0} = 100 \text{ G}$ ,  $R_i = 1 \text{ m}$  and  $T_e = 10 \text{ eV}$ . This estimation means the change of the radial electric field is the same order of the perturbed magnetic field to the unperturbed magnetic field.




Fig. 1. Angular equilibrium frequency of electrons, which is hollow distribution with the inner and outer radius are 2.0 and 3.0, respectively.